

Health impact assessment of PM₁₀ exposures in the city of Caen, France.

Glennec P., Monroux F. Ecole Nationale de la Santé Publique (National School of Public Health), Rennes, France.

Context and objectives

The city of Caen is located in north-western France. Air pollution is mainly due to traffic sources. The aim of this work is to assess the public health impact of both short (with short term effects) and chronic (with long term effects) exposures to PM₁₀.

Methods

The standard World Health Organisation (WHO) methodology (1) for an Health Impact Assessment (HIA) has been used to calculate the **attributable deaths and hospital admissions**. Population exposure is estimated from PM₁₀ (Particulate Matter <10 µm) concentrations collected by the local air quality measurement network : Aircom. The relative risks have been modelled by the **exposure-risk functions** established in epidemiologic studies carried out in the general population. The Aphea-2 (2-4) program, which combines european time-series studies, have been used for short exposures effects, and a meta risk calculated (5) from cohort studies for long exposures effects. Health impact have been calculated with the EIS-PA software (6) to assess :

- for the short exposures : total impact with reference to the baseline level of 10 µg/m³, health gain for a daily 10 % reduction of pollution, health gain for a daily compliance to 40 µg/m³ (EU regulatory level for annual mean in 2005)
- for the chronic exposures : health gain for a 5 µg/m³ annual reduction.




Fig 1. HIA principle

| | PM10 (µg/m ³) |
|--------------------|---------------------------|
| Min | 2 |
| P 5 | 7 |
| P 25 | 12 |
| P 50 | 17 |
| P 75 | 25 |
| P 95 | 42 |
| Max | 87 |
| Daily mean | 19 |
| Standard deviation | 11,4 |
| % missing values | 1% |

Table 1. PM 10 outdoor exposures, Caen 1998-2002.


Findings

Table 2 : health impact of outdoor particulate air pollution. Caen 1998-2002.

| Health effect | Attributable cases (CI 95 %) |
|--|------------------------------|
| Anticipated death | 26 (17-35) |
| Short term Respiratory hospital admissions | 16 (11-23) |
| Cardiovascular hospital admissions | 27 (11-44) |
| Long term Mortality | 90 (54-126) * |


* for a 5 µg/m³ reduction of the annual mean.

Fig 2. Health efficiency of particulate air pollution policy scenarios, Caen 1998-2002.



Reducing daily levels of pollution is more efficient than avoiding peaks. Health impact is mainly due to moderate, but frequent, levels of pollution.

Fig 3. Health impact vs exposure levels. Caen 1998-2000.



Discussion

Some major uncertainties and ways of reducing it :

- long term effects : validity in Europe, and accuracy of relative risks : to go ahead with European cohort studies

- hazard identification/health end points : major effects such as bronchitis and asthma attacks could not be assessed : to know the local baseline risks in France

Variability and its uses :

- geographic : when necessary (town planning), needs to be assessed with population time activity patterns inquiries and city wide PM10 measurements

- temporal : it has been used here to describe the respective contribution of polluted and 'normal' days.

References

1. WHO Regional Office for Europe. Quantification of the health effects of exposure to air pollution. Report on a WHO working group, Bilthoven, The Netherlands, November 2000; 29 pages.
2. Katsouyanni K, et al. Confounding and effect modification in the short-term effects of ambient particles on total mortality: results from 29 European cities within the APHEA2 project. *Epidemiology*. 2001 Sep;12(5):521-31.
3. Atkinson RW et al. Acute effects of particulate air pollution on respiratory admissions : results from APHEA 2 project. *Air Pollution and Health: a European Approach. Am J Respir Crit Care Med.*, 2001 Nov 15;164(10 Pt 1):1860-6.
4. Le Tertre A, et al. Short-term effects of particulate air pollution on cardiovascular diseases in eight European cities. *J Epidemiol Community Health*, 2002 Oct;56(10):773-9.
5. Kunzli N, et al. Public-health impact of outdoor and traffic-related air pollution: a European assessment. *Lancet* 2000; 356:795-80.
6. Institut de Veille Sanitaire. Évaluation d'impact sanitaire de la pollution atmosphérique urbaine. Actualisation du guide méthodologique 2003. St Maurice, France, 2003; 32 pages.

